arXiv:1006.1140v2 [math.CA] 14 Dec 2010

Nonsymmetric Askey-Wilson polynomials as vector-valued polynomials

Tom H. Koornwinder* and Fethi Bouzeffour†

Dedicated to Paul Butzer on the occasion of his 80th birthday

Abstract

Nonsymmetric Askey-Wilson polynomials are usually written as Laurent polynomials. We write them equivalently as 2-vector-valued symmetric Laurent polynomials. Then the Dunkl-Cherednik operator of which they are eigenfunctions, is represented as a 2×2 matrix-valued operator. As a new result made possible by this approach we obtain positive definiteness of the inner product in the orthogonality relations, under certain constraints on the parameters. A limit transition to nonsymmetric little q-Jacobi polynomials also becomes possible in this way. Nonsymmetric Jacobi polynomials are considered as limits both of the Askey-Wilson and of the little q-Jacobi case.

1 Introduction

Originally, most orthogonal special functions associated with root systems, like the Heckman-Opdam Jacobi polynomials and the Macdonald polynomials, were Weyl group invariant functions. By work by Dunkl, Heckman and in particular Cherednik it was next shown that there are related orthogonal systems of special functions which are not Weyl group invariant, but are in a sense more simple, and from which the earlier Weyl group invariant special functions can be obtained by symmetrization. While Cherednik's theory, involving double affine Hecke algebras, has been developed for general root systems, to a large extent independent of the classification, the specialization of this theory to the case of rank one has its own interest, because everything can be done there in a much more explicit way, and new results for special functions in one variable can be obtained. In the rank one case the Weyl group has order 2, and Weyl group symmetry turns down to a symmetry under the map $t\mapsto -t$ or $z\mapsto z^{-1}$.

The case of nonsymmetric Askey-Wilson polynomials was first considered briefly by Sahi [19], [20] as a specialization for $l=1$ of his extension of Cherednik's theory to the pair of root systems $(C_l^\vee, C_l)$. The case $l=1$ was treated afterwards in more detail by Noumi and Stokman

*Korteweg-de Vries Institute, University of Amsterdam, P.O. Box 94248, 1090 GE Amsterdam, Netherlands; T.H.Koornwinder@uva.nl

†Institut Préparatoire aux Études d'Ingénieur de Bizerte, 8000 Bizerte, Tunisia; bouzeffourfethi@yahoo.fr

[16] and by Macdonald [14, §6.6]. See also the first author's paper [13]. In all these references the nonsymmetric Askey-Wilson polynomials are defined as certain Laurent polynomials and the bilinear form for which they are orthogonal (or rather biorthogonal) is given in terms of a contour integral.

In the present paper we give a presentation of nonsymmetric Askey-Wilson polynomials as two-dimensional vector-valued polynomials. The vector is a pair of symmetric Laurent polynomials: an Askey-Wilson polynomial and another one with the degree lowered by 1 and two parameters raised by 1. As a first application of this approach we can treat the orthogonality in the nonsymmetric case by reducing it to the well-known orthogonality of the two components of the vector. We obtain the new result that, under certain constraints on the parameters, there is orthogonality with respect to a positive definite inner product. It does not seem possible to read off this result from a positive weight function in the contour integral.

As a second application we can easier take limits of nonsymmetric Askewy-Wilson polynomials corresponding to the familiar limits of the symmetric polynomials in the (q-)Askey scheme. In this paper we consider the limit from nonsymmetric Askey-Wilson polynomials to nonsymmetric little q-Jacobi polynomials. Again, the vector-valued approach is crucial here. Next we obtain nonsymmetric Jacobi polynomials as limits in two different ways: directly as a limit of the Askey-Wilson case, where the Laurent polynomials can be used, and as a limit of the little q-Jacobi case, where we need the vector-valued polynomials. We start the exposition by recalling the well-known nonsymmetric Bessel case, and we end by obtaining nonsymmetric Bessel as a limit of nonsymmetric Jacobi.

We expect that to every limit arrow in the (q-)Askey scheme there corresponds a limit transition in the nonsymmetric case, and also for limits going out of the (q-)Askey scheme to various (q-)Bessel functions. We hope to present some further examples in a subsequent paper. Another topic which we will not yet treat here, is a study of the corresponding limit algebras of the double affine Hecke algebra in the Askey-Wilson case.

Conventions Throughout assume $0 < q < 1$. For (q-)Pochhammer symbols and (q-)hypergeometric series use the notation of [7]. In particular, for $n \geq 0$,

$$(a;q)_n := \prod_{j=0}^{n-1}(1-aq^j), \qquad (a_1,\dots,a_r;q)_n := (a_1;q)_n \cdots (a_r;q)_n\,,$$

$${}_r\phi_{r-1}\left(\begin{matrix} q^{-n}, a_2, \dots, a_r \\ b_1, \dots, b_{r-1}\end{matrix}; q, z\right) := \sum_{k=0}^{n} \frac{(q^{-n}, a_2, \dots, a_r; q)_k}{(b_1, \dots, b_{r-1}, q; q)_k}\, z^k.$$

Let e_1, e_2, e_3, e_4 be the elementary symmetric polynomials in a, b, c, d:

$$e_1 := a+b+c+d, \qquad e_2 := ab+ac+bc+ad+bd+cd,$$
$$e_3 := abc+abd+acd+bcd, \qquad e_4 := abcd. \tag{1.1}$$

For Laurent polynomials f in z the z-dependence will be written as $f[z]$. Symmetric Laurent

polynomials $f[z] = \sum_{k=-n}^{n} c_k z^k$ (where $c_k = c_{-k}$) are related to ordinary polynomials $f(x)$ in $x = \frac{1}{2}(z + z^{-1})$ by $f(\frac{1}{2}(z + z^{-1})) = f[z]$.

2 The nonsymmetric Hankel transform

For Bessel functions J_α see [15, Ch. 10] and references given there. We will work with differently normalized Bessel functions

$$\mathcal{J}_\alpha(x) := \Gamma(\alpha + 1)\,(2/x)^\alpha\, J_\alpha(x). \tag{2.1}$$

Then (see [15, (10.16.9)])

$$\mathcal{J}_\alpha(x) = \sum_{k=0}^{\infty} \frac{(-\frac{1}{4}x^2)^k}{(\alpha+1)_k\, k!} = {}_0F_1\left(\begin{matrix} - \\ \alpha + 1 \end{matrix}; -\tfrac{1}{4}x^2\right) \qquad (\alpha > -1).$$

$\mathcal{J}_\alpha$ is an entire analytic function and we have the simple properties and special cases

$$\mathcal{J}_\alpha(x) = \mathcal{J}_\alpha(-x), \quad \mathcal{J}_\alpha(0) = 1, \quad \mathcal{J}_{-1/2}(x) = \cos x, \quad \mathcal{J}_{1/2}(x) = \frac{\sin x}{x}\,.$$

The functions $x \mapsto \mathcal{J}_\alpha(\lambda x)$ satisfy the eigenvalue equation [15, (10.13.5)]:

$$\left(\frac{d^2}{dx^2} + \frac{2\alpha+1}{x}\,\frac{d}{dx}\right)\mathcal{J}_\alpha(\lambda x) = -\lambda^2\,\mathcal{J}_\alpha(\lambda x).$$

The *Hankel transform* pair [15, §10.22(v)], for f in a suitable function class, is given by

$$\begin{cases} \widehat{f}(\lambda) = \displaystyle\int_0^\infty f(x)\mathcal{J}_\alpha(\lambda x) x^{2\alpha+1}\,dx, \\ f(x) = \dfrac{1}{2^{2\alpha+1}\Gamma(\alpha+1)^2}\displaystyle\int_0^\infty \widehat{f}(\lambda)\mathcal{J}_\alpha(\lambda x)\lambda^{2\alpha+1}\,d\lambda. \end{cases} \tag{2.2}$$

Now consider the so-called *nonsymmetric Bessel function*, also called *Dunkl-type Bessel function*, in the rank one case (see [5, §4], [18, Example 2.29]):

$$\mathcal{E}_\alpha(x) := \mathcal{J}_\alpha(x) + \frac{i\,x}{2(\alpha+1)}\,\mathcal{J}_{\alpha+1}(x). \tag{2.3}$$

In particular, $\mathcal{E}_{-1/2}(x) = e^{ix}$. The nonsymmetric Hankel transform pair takes the form

$$\begin{cases} \widehat{f}(\lambda) = \displaystyle\int_{-\infty}^\infty f(x)\,\mathcal{E}_\alpha(-\lambda x)\,|x|^{2\alpha+1}\,dx, \\ f(x) = \dfrac{1}{2^{2(\alpha+1)}\Gamma(\alpha+1)^2}\displaystyle\int_{-\infty}^\infty \widehat{f}(\lambda)\,\mathcal{E}_\alpha(\lambda x)\,|\lambda|^{2\alpha+1}\,d\lambda. \end{cases} \tag{2.4}$$

The transform pair (2.4) follows immediately from (2.2) by putting $f(x) = f_1(x) + xf_2(x)$ in (2.4) with f_1 and f_2 even. For given α define the differential-reflection operator

$$(Yf)(x) := f'(x) + (\alpha + \tfrac{1}{2})\,\frac{f(x) - f(-x)}{x}\,. \tag{2.5}$$

This is the *Dunkl operator* for root system A_1 (see [6, Definition 4.4.2)]). Then we have the eigenvalue equation

$$Y(\mathcal{E}_\alpha(\lambda\,.\,)) = i\,\lambda\,\mathcal{E}_\alpha(\lambda\,.\,)\,. \tag{2.6}$$

If, in (2.6), we substitute (2.3), compare even and odd parts, and then substitute (2.1), then we see that (2.6) is equivalent with a pair of lowering and raising differentiation formulas for Bessel functions (see [15, (10.6.2)]):

$$J'_\alpha(x) - \frac{\alpha}{x}\,J_\alpha(x) = -J_{\alpha+1}(x), \qquad J'_{\alpha+1}(x) + \frac{\alpha+1}{x}\,J_{\alpha+1}(x) = J_\alpha(x).$$

3 Askey-Wilson polynomials

Askey-Wilson polynomials were introduced in [1], see also [7, §7.5] and [10, §3.1]. We will consider these polynomials in $x = (z + z^{-1})/2$ as symmetric Laurent polynomials in z and we will renormalize them such that they are *monic*, i.e., the coefficient of the highest degree term in z is 1:

$$\begin{aligned} P_n[z] = P_n[z; a, b, c, d \mid q] &= P_n\big(\tfrac{1}{2}(z + z^{-1})\big) \\ &:= \frac{(ab, ac, ad; q)_n}{a^n (abcdq^{n-1}; q)_n}\;{}_4\phi_3\left(\begin{matrix} q^{-n}, q^{n-1}abcd, az, az^{-1} \\ ab, ac, ad \end{matrix}; q, q\right). \end{aligned} \tag{3.1}$$

They are eigenfunctions of a second order q-difference operator $L = L_{a,b,c,d;q}$:

$$(LP_n)[z] = (q^{-n} - 1)(1 - abcdq^{n-1})P_n[z], \tag{3.2}$$

where

$$(Lf)[z] := A[z]\,f[qz] + A[z^{-1}]\,f[q^{-1}z] - \big(A[z] + A[z^{-1}]\big)\,f[z]$$

and

$$A[z] := \frac{(1 - az)(1 - bz)(1 - cz)(1 - dz)}{(1 - z^2)(1 - qz^2)}\,.$$

The Askey-Wilson polynomials are on the top level of the q-Askey scheme (see [10, §3]; a graphical display of this scheme including the arrows will appear in [11], just before Chapter 14).

The three-term recurrence relation for the polynomials (3.1) is as follows (see [10, (3.1.5)]):

$$(z + z^{-1})P_n[z] = P_{n+1}[z] + B_n P_n[z] + C_n P_{n-1}[z], \tag{3.3}$$

where $C_nP_{n-1}[z] := 0$ for $n = 0$, and where B_n and C_n (partially expresed in terms of the e_i, see (1.1)) are given by:

$$B_n := q^{n-1}\,\frac{(1-q^n-q^{n+1})e_3+qe_1+q^{2n-1}e_3e_4-q^{n-1}(1+q-q^{n+1})e_1e_4}{(1-q^{2n-2}e_4)(1-q^{2n}e_4)}\,, \tag{3.4}$$

$$\begin{aligned} C_n := {} & (1-q^{n-1}ab)(1-q^{n-1}ac)(1-q^{n-1}ad)(1-q^{n-1}bc)(1-q^{n-1}bd)(1-q^{n-1}cd) \\ & \times \frac{(1-q^n)(1-q^{n-2}e_4)}{(1-q^{2n-3}e_4)(1-q^{2n-2}e_4)^2(1-q^{2n-1}e_4)}\,. \end{aligned} \tag{3.5}$$

From (3.4) and (3.5) it is clear that $P_n[z;a,b,c,d \mid q]$ is symmetric in a,b,c,d (well-known, but not yet evident from (3.1)).

By Favard's theorem (see [3, Theorems I.4.4 and II.3.2]), there exists a positive Borel measure $\mu = \mu_{a,b,c,d;q}$ on $\mathbb{R}$ with $\mu(\mathbb{R}) = 1$ such that

$$\langle P_m, P_n\rangle = \langle P_m, P_n\rangle_{a,b,c,d;q} := \int_{\mathbb{R}} P_m(x)\,P_n(x)\,d\mu(x) = h_n\,\delta_{m,n} \tag{3.6}$$

with

$$h_n = h_n^{a,b,c,d;q} = C_1C_2\ldots C_n = \frac{(q,ab,ac,ad,bc,bd,cd;q)_n}{(abcd;q)_{2n}(q^{n-1}abcd;q)_n} > 0. \tag{3.7}$$

if and only if

$$B_n \text{ is real for } n \geq 0 \text{ and } C_n > 0 \text{ for } n \geq 1. \tag{3.8}$$

A sufficient condition for (3.8) to hold is that among a,b,c,d there are two, one or zero pairs of complex conjugates with the other parameters being real and that pairwise products of parameters are less than 1 in absolute value. Then the orthogonality measure μ can be given explicitly (see [1, §2], [10, §3.1]).

4 Nonsymmetric Askey-Wilson polynomials

From now on also assume:

$$a,b,c,d \neq 0, \quad abcd \neq q^{-m}(m = 0,1,2,\ldots), \quad \{a,b\} \cap \{a^{-1},b^{-1}\} = \emptyset. \tag{4.1}$$

In terms of

$$\begin{aligned} P_n[z] \; &= P_n[z;a,b,c,d \mid q], \\ Q_n[z] &:= a^{-1}b^{-1}z^{-1}(1-az)(1-bz)\,P_{n-1}[z;qa,qb,c,d \mid q] \end{aligned}$$

we define the *nonsymmetric Askey-Wilson polynomials* by:

$$E_{-n} := P_n - Q_n \qquad (n = 1,2,\ldots), \tag{4.2}$$

$$E_n := P_n - \frac{ab(1-q^n)(1-q^{n-1}cd)}{(1-q^nab)(1-q^{n-1}abcd)}\,Q_n \qquad (n = 0,1,2,\ldots), \tag{4.3}$$

with the convention that $(1-q^n)Q_n := 0$ for $n = 0$. They are eigenfunctions of a q-difference-reflection operator

$$\begin{aligned}(Yf)[z] := q^{-1}abcd\, f[z] &+ \frac{(1-az)(1-bz)(1-cz)(1-dz)}{(1-z^2)(1-qz^2)}(f[qz]-f[z])\\ &+\frac{(1-az)(1-bz)\big((c+d)qz-(cd+q)\big)}{q(1-z^2)(1-qz^2)}\,(f[z^{-1}]-f[z])\\ &+\frac{(c-z)(d-z)\big(1+ab-(a+b)z\big)}{(1-z^2)(q-z^2)}\,(f[qz^{-1}]-f[z]).\end{aligned} \tag{4.4}$$

Then

$$YE_{-n} = q^{-n}\,E_{-n} \qquad (n = 1, 2, \ldots), \tag{4.5}$$

$$YE_n = q^{n-1}abcd\,E_n \qquad (n = 0, 1, 2, \ldots). \tag{4.6}$$

The idea of working with these functions E_n and operator Y comes from Cherednik's theory [2] of double affine Hecke algebras associated with root systems, extended by S. Sahi [19], [20] to the type $(C_l^\vee, C_l)$. The special case $l = 1$ was treated afterwards in [16], [14, §6.6] and [13].

The *double affine Hecke algebra* of type $(C_1^\vee, C_1)$ is the algebra $\tilde{\mathfrak{H}}$ generated by Z, Z^{-1}, T_1, T_0 with relations $ZZ^{-1} = 1 = Z^{-1}Z$ and

$$(T_1 + ab)(T_1 + 1) = 0, \qquad (T_0 + q^{-1}cd)(T_0 + 1) = 0,$$
$$(T_1Z + a)(T_1Z + b) = 0, \qquad (qT_0Z^{-1} + c)(qT_0Z^{-1} + d) = 0.$$

This algebra acts faithfully on the space of Laurent polynomials:

$$\begin{aligned}(Zf)[z] &:= z\,f[z],\\ (T_1f)[z] &:= \frac{(a+b)z-(1+ab)}{1-z^2}\,f[z] + \frac{(1-az)(1-bz)}{1-z^2}\,f[z^{-1}],\\ (T_0f)[z] &:= \frac{q^{-1}z((cd+q)z-(c+d)q)}{q-z^2}\,f[z] - \frac{(c-z)(d-z)}{q-z^2}\,f[qz^{-1}].\end{aligned}$$

Then $Y = T_1T_0$ acts on the space of symmetric Laurent polynomials as (4.4).

The following Proposition (see [13, Proposition 3.1]) will enable us to rewrite (4.5) and (4.6) in vector-valued form.

Proposition 4.1.
(a) *T_1 acting on Laurent polynomials has eigenvalues $-ab$ and -1.*
(b) *$T_1f = -ab\,f \iff f$ is symmetric.*
(c) *$T_1f = -f \iff f[z] = z^{-1}(1-az)(1-bz)g[z]$ for some symmetric Laurent polynomial g.*

Let A be an operator acting on the space of Laurent polynomials. Write

$$f[z] = f_1[z] + z^{-1}(1-az)(1-bz)f_2[z] \quad (f_1, f_2 \text{ symmetric Laurent polynomials}). \tag{4.7}$$

Then we can write

$$(Af)[z] = (A_{11}f_1 + A_{12}f_2)[z] + z^{-1}(1-az)(1-bz)(A_{21}f_1 + A_{22}f_2)[z], \tag{4.8}$$

where the $A_{i\,j}$ are operators acting on the space of symmetric Laurent polynomials. So we have the identifications

$$f \leftrightarrow \begin{pmatrix} f_1 \\ f_2 \end{pmatrix}, \quad A \leftrightarrow \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}. \tag{4.9}$$

In particular, we have

$$E_{-n}[z] = \begin{pmatrix} P_n[z;a,b,c,d \mid q] \\ -a^{-1}b^{-1}P_{n-1}[z;qa,qb,c,d \mid q] \end{pmatrix} \quad (n = 1,2,\ldots), \tag{4.10}$$

$$E_n[z] = \begin{pmatrix} P_n[z;a,b,c,d \mid q] \\ -\dfrac{(1-q^n)(1-q^{n-1}cd)}{(1-q^nab)(1-q^{n-1}abcd)}\, P_{n-1}[z;qa,qb,c,d \mid q] \end{pmatrix} \quad (n = 0,1,2,\ldots), \tag{4.11}$$

where $(1-q^n)P_{n-1} := 0$ for $n = 0$. Also $Y = \begin{pmatrix} Y_{11} & Y_{12} \\ Y_{21} & Y_{22} \end{pmatrix}$ with

$$Y_{11} = q^{-1}abcd - \frac{ab}{1-ab}\, L_{a,b,c,d;q}\,, \tag{4.12}$$

$$Y_{22} = \frac{1 - abcd - abq + abcdq + L_{aq,bq,c,d;q}}{q(1-ab)} \tag{4.13}$$

(recall that $L_{a,b,c,d;q}$ is the second order q-difference operator occurring in (3.2)) and

$$(Y_{21}g)[z] = \frac{z(c-z)(d-z)\left(g[q^{-1}z] - g[z]\right)}{(1-ab)(1-z^2)(1-qz^2)} + \frac{z(1-cz)(1-dz)\left(g[qz] - g[z]\right)}{(1-ab)(1-z^2)(1-qz^2)}\,, \tag{4.14}$$

$$\begin{aligned}(Y_{12}h)[z] = {} & \frac{ab(a-z)(b-z)(1-az)(1-bz)}{(1-ab)z(q-z^2)(1-qz^2)}\left((cd+q)(1+z^2) - (1+q)(c+d)z\right)h[z] \\ & - \frac{ab(a-z)(b-z)(c-z)(d-z)(aq-z)(bq-z)}{q(1-ab)z(1-z^2)(q-z^2)}\, h[q^{-1}z] \\ & - \frac{ab(1-az)(1-bz)(1-cz)(1-dz)(1-aqz)(1-bqz)}{q(1-ab)z(1-z^2)(1-qz^2)}\, h[qz]. \end{aligned} \tag{4.15}$$

The eigenvalue equations (4.5), (4.6) for E_n and for E_{-n} are equivalent to the four equations

$$L_{a,b,c,d;q}P_n[\,.\,;a,b,c,d\mid q] = (q^{-n}-1)(1-abcdq^{n-1})P_n[\,.\,;a,b,c,d\mid q],$$
$$L_{qa,qb,c,d;q}P_{n-1}[\,.\,;qa,qb,c,d\mid q] = (q^{-n+1}-1)(1-abcdq^{n})P_{n-1}[\,.\,;qa,qb,c,d\mid q],$$
$$Y_{21}P_n[\,.\,;a,b,c,d\mid q] = -\frac{(q^{-n}-1)(1-cdq^{n-1})}{1-ab}\,P_{n-1}[\,.\,;qa,qb,c,d\mid q],$$
$$Y_{12}P_{n-1}[\,.\,;qa,qb,c,d\mid q] = -\frac{ab(q^{-n}-ab)(1-abcdq^{n-1})}{1-ab}\,P_n[\,.\,;a,b,c,d\mid q].$$

5 Orthogonality relations

Consider $\langle\,.\,,\,.\,\rangle_{a,b,c,d;q}$ (see (3.6), (3.7)) as a symmetric bilinear form on the space of symmetric Laurent polynomials. With the identification $f \leftrightarrow (f_1, f_2)$ between a Laurent polynomial f and a pair of symmetric Laurent polynomials (see (4.7)) we look for a symmetric bilinear form on the space of Laurent polynomials of the form

$$\langle g,h\rangle = \langle (g_1,g_2),(h_1,h_2)\rangle = \langle g_1,h_1\rangle_{a,b,c,d;q} + C\langle g_2,h_2\rangle_{qa,qb,c,d;q} \tag{5.1}$$

such that the nonsymmetric Askey-Wilson polynomials E_n ($n\in\mathbb{Z}$) given by (4.2), (4.3) are orthogonal with respect to this form, i.e.

$$\langle E_m,E_n\rangle = 0 \quad (m\neq n). \tag{5.2}$$

By (4.10), (4.11) the orthogonality certainly holds if $|n|\neq|m|$. Thus we have to determine C in (5.1) such that $\langle E_n,E_{-n}\rangle = 0$. By (4.2), (4.3) this turns down to

$$C = -ab\,\frac{(1-q^nab)(1-q^{n-1}abcd)}{(1-q^n)(1-q^{n-1}cd)}\,\frac{h_n^{a,b,c,d;q}}{h_{n-1}^{qa,qb,c,d;q}}\,.$$

A priori, it is not clear that C is independent of n. However, from (3.7) we compute

$$\frac{h_n^{a,b,c,d;q}}{h_{n-1}^{qa,qb,c,d;q}} = \frac{(1-q^n)(1-q^{n-1}cd)}{(1-q^nab)(1-q^{n-1}abcd)}\,\frac{(1-ab)(1-qab)(1-ac)(1-ad)(1-bc)(1-bd)}{(1-abcd)(1-qabcd)}\,.$$

Thus C is independent of n and the form (5.1) becomes more explicitly

$$\langle g,h\rangle = \langle g_1,h_1\rangle_{a,b,c,d;q} - ab\,\frac{(1-ab)(1-qab)(1-ac)(1-ad)(1-bc)(1-bd)}{(1-abcd)(1-qabcd)}\,\langle g_2,h_2\rangle_{qa,qb,c,d;q}\,. \tag{5.3}$$

With respect to the form (5.3) we have thus shown that the orthogonality (5.2) holds.

Under some further assumptions we can even show that (5.3) defines a positive definite inner product.

Proposition 5.1. *Let $a,b,c,d;q$, besides satisfying* (4.1)*, also be such that condition* (3.8) *holds both for $a,b,c,d;q$ and for $qa,qb,c,d;q$. Moreover assume that $ab<0$. Then $cd<1$, all coefficients in* (4.2), (4.3) *are real, and the inner product defined by* (5.3) *is positive definite.*

Proof Since C_n for $a,b,c,d;q$ and C_{n-1} for $qa,qb,c,d;q$ are positive, their quotient must be positive. Hence, by (3.5),

$$\frac{(1-q^{n-1}cd)}{(1-q^{n-2}cd)}\,\frac{(1-q^{n-2}abcd)}{(1-q^{n-1}abcd)}>0 \qquad (n\geq 2).$$

Taking the product of these inequalities and telescoping yields $\frac{1-abcd}{1-cd}>0$. Since $ab<0$, we conclude that cd is real. Then $cd\geq 1$ is impossible, so $cd<1$. Hence $abcd<1$. Then we also see that all coefficients in (4.2), (4.3) are real. Finally, since

$$C_1=\frac{(1-ab)(1-ac)(1-ad)(1-bc)(1-bd)(1-cd)(1-q)}{(1-abcd)^2(1-qabcd)}>0,$$

we have

$$C=-ab\,\frac{(1-ab)(1-qab)(1-ac)(1-ad)(1-bc)(1-bd)}{(1-abcd)(1-qabcd)}>0,$$

and the positive definiteness of (5.3) is settled. □

In Noumi & Stokman [16, Proposition 6.8] a biorthogonality result involving the system of Laurent polynomials E_n is given with respect to a bilinear form defined in terms of a contour integral. This is closely connected with our orthogonality (5.2). Our positive definiteness result is probably new. It does not seem to be possible to rewrite the contour integral in [16] such that a positive definite hermitian inner product will become apparent under suitable constraints on the parameters. In the $q=1$ limit case this will turn out to be much nicer, see (7.19).

6 From nonsymmetric Askey-Wilson to nonsymmetric little q-Jacobi

We consider *little q-Jacobi polynomials* (see [10, §3.12]) in monic form:

$$P_n(x;a,b;q):=\frac{(-1)^n q^{n(n-1)/2}(aq;q)_n}{(abq^{n+1};q)_n}\;{}_2\phi_1\left(\begin{matrix}q^n,abq^{n+1}\\ aq\end{matrix};q,qx\right). \tag{6.1}$$

They are limits of Askey-Wilson polynomials (3.1) (see [12, Proposition 6.3]):

$$P_n(x;a,b;q)=\lim_{\lambda\downarrow 0}\lambda^n P_n[\lambda^{-1}x;-q^{1/2}a,qb\lambda,-q^{1/2},\lambda^{-1}\mid q]. \tag{6.2}$$

Note that we take the limit of symmetric Laurent polynomials in x, but in the limit we have ordinary polynomials in x, since all negative powers of x are killed in the limit.

The polynomials (6.1) are eigenfunctions of a second order q-difference operator $L = L_{a,b;q}$:

$$(LP_n)(x) = (q^{-n} - 1)(1 - abq^{n+1})P_n(x), \tag{6.3}$$

where

$$(Lf)(x) := A(x)\, f(qx) + B(x)\, f(q^{-1}x) - \big(A(x) + B(x)\big)\, f(x)$$

and

$$A(x) := \frac{abqx - a}{x}\,, \qquad B(x) := \frac{x-1}{x}\,.$$

The system of polynomials (6.1) is orthogonal with respect to a positive orthogonality measure if $0 < a < q^{-1}$ and $b < q^{-1}$. This can be given as an explicit q-integral, see [10, (3.12.2)]. In the orthogonality relations

$$\langle P_m, P_n\rangle = \langle P_m, P_n\rangle_{a,b;q} = h_n\, \delta_{m,n}$$

we have

$$h_n = h_n^{a,b;q} = \frac{q^{n^2} a^n (q, aq, bq; q)_n}{(abq^2; q)_{2n} (abq^{n+1}; q)_n}\,.$$

Corresponding to (6.2) there are limits of nonsymmetric Askey-Wilson polynomials (4.10), (4.11) (in vector-valued form) which yield *nonsymmetric little q-Jacobi polynomials* in vector-valued form:

$$E_n(x; a, b, q) := \lim_{\lambda \downarrow 0} \lambda^n E_n[\lambda^{-1}x; -q^{1/2}a, qb\lambda, -q^{1/2}, \lambda^{-1} \mid q] \qquad (n \in \mathbb{Z}).$$

Their expressions in vector-valued form (with the usual convention for $n = 0$) are

$$E_{-n}(x; a, b, q) = \begin{pmatrix} P_n(x; a, b; q) \\ q^{-3/2} a^{-1} b^{-1} P_{n-1}(x; qa, qb; q) \end{pmatrix} \quad (n = 1, 2, \ldots), \tag{6.4}$$

$$E_n(x; a, b, q) = \begin{pmatrix} P_n(x; a, b; q) \\ -\dfrac{q^{n-\frac{1}{2}}(1 - q^n)}{1 - q^{n+1}ab}\, P_{n-1}(x; qa, qb; q) \end{pmatrix} \quad (n = 0, 1, 2, \ldots). \tag{6.5}$$

Note that taking corresponding limits of the nonsymmetric Askey-Wilson polynomials (4.2), (4.3) (as Laurent polynomials) would give a system of linearly dependent ordinary polynomials.

The 2×2 matrix-valued operator $Y = \begin{pmatrix} Y_{11} & Y_{12} \\ Y_{21} & Y_{22} \end{pmatrix}$ with entries given by (4.12)–(4.15) also has a limit for $\lambda \downarrow 0$ after the rescaling $a \to -q^{1/2}a$, $b \to qb\lambda$, $c \to -q^{1/2}$, $d \to \lambda^{-1}$, $z \to \lambda^{-1}x$. Thus, if the eigenvalue equations (4.5), (4.6) are rescaled in this way with both sides being

multiplied by λ^n, then in the limit for $\lambda \downarrow 0$ we obtain

$$\left(\begin{pmatrix} Y_{11} & Y_{12} \\ Y_{21} & Y_{22} \end{pmatrix} - q^{-n}\right) \begin{pmatrix} P_n(x;a,b;q) \\ q^{-3/2}a^{-1}b^{-1}P_{n-1}(x;qa,qb;q) \end{pmatrix} = 0 \qquad (n>0), \tag{6.6}$$

$$\left(\begin{pmatrix} Y_{11} & Y_{12} \\ Y_{21} & Y_{22} \end{pmatrix} - q^{n+1}ab\right) \begin{pmatrix} P_n(x;a,b;q) \\ -\dfrac{q^{n-\frac{1}{2}}(1-q^n)}{1-q^{n+1}ab}\, P_{n-1}(x;qa,qb;q) \end{pmatrix} = 0 \qquad (n\geq 0). \tag{6.7}$$

Here

$$\begin{aligned} &Y_{11} = qab, \qquad Y_{22} = q^{-1} - q(1-q)ab + q^{-1}L_{aq,bq;q}, \\ &(Y_{21}g)(x) = \frac{g(x)-g(qx)}{q^{1/2}x}, \quad (Y_{12}g)(x) = a^2bq^{3/2}(1-bqx)g(x) - abq^{1/2}(1-x)g(q^{-1}x). \end{aligned} \tag{6.8}$$

In a similar way as in section 5 we can prove that that the vector-valued polynomials E_n are orthogonal with respect to the symmetric bilinear form

$$\langle g,h\rangle = \langle (g_1,g_2),(h_1,h_2)\rangle = \langle g_1,h_1\rangle_{a,b;q} + \frac{q^2a^2b(1-qa)(1-qb)}{(1-q^2ab)(1-q^3ab)}\,\langle g_2,h_2\rangle_{qa,qb;q}\,. \tag{6.9}$$

This form is positive definite if $b>0$, i.e., if $a,b\in(0,q^{-1})$.

7 Limits to nonsymmetric Jacobi polynomials

Consider *Jacobi polynomials* (see [10, §1.8]) as monic symmetric Laurent polynomials:

$$\begin{aligned} P_n[z;\alpha,\beta] = P_n\big((z+z^{-1})/2;\alpha,\beta\big) &:= \frac{2^{2n}(\alpha+1)_n}{(n+\alpha+\beta+1)_n}\ {}_2F_1\left(\begin{matrix} -n, n+\alpha+\beta+1 \\ \alpha+1 \end{matrix}; \frac{2-z-z^{-1}}{4}\right) \\ &= \frac{2^{2n}n!}{(n+\alpha+\beta+1)_n}\,P_n^{(\alpha,\beta)}\big((z+z^{-1})/2\big). \end{aligned} \tag{7.1}$$

For $\alpha,\beta>-1$ these polynomials satisfy the orthogonality relations

$$\langle P_m,P_n\rangle = \langle P_m,P_n\rangle_{\alpha,\beta} = \frac{2^{-(\alpha+\beta+1)}\Gamma(\alpha+\beta+2)}{\Gamma(\alpha+1)\Gamma(\beta+1)}\int_{-1}^{1} P_m(x)\,P_n(x)\,(1-x)^\alpha(1+x)^\beta\,dx = h_n\,\delta_{m,n}, \tag{7.2}$$

where

$$h_n = h_n^{\alpha,\beta} = \frac{2^{4n}(\alpha+1)_n(\beta+1)_n n!}{(\alpha+\beta+2)_{2n}(n+\alpha+\beta+1)_n}\,.$$

Jacobi polynomials are limits of Askey-Wilson polynomials, or rather of continuous q-Jacobi polynomials, a two-parameter subclass of the Askey-Wilson polynomials (see [10, (3.10.14), (5.10.2)]):

$$P_n[z;\alpha,\beta] = \lim_{q\uparrow 1} P_n\big[z;q^{\alpha+\frac{1}{2}},-q^{\beta+\frac{1}{2}},q^{\frac{1}{2}},-q^{\frac{1}{2}}\mid q\big]. \tag{7.3}$$

The ordering of parameters in the right-hand side of (7.3), different from the ordering in [10, (3.10.14)] but allowed in view of the symmetry in the parameters, was chosen in order to be able to take limits of the formulas in section 4. Indeed, we can now obtain nonsymmetric Jacobi polynomials as limits

$$E_n[z;\alpha,\beta] := \lim_{q\uparrow 1} E_n\big[z; q^{\alpha+\frac{1}{2}}, -q^{\beta+\frac{1}{2}}, q^{\frac{1}{2}}, -q^{\frac{1}{2}} \mid q\big]$$

of nonsymmetric Askey-Wilson polynomials (4.2), (4.3) by using (7.3). Then

$$E_{-n}[z;\alpha,\beta] = P_n[z;\alpha,\beta] - (z-z^{-1})\,P_{n-1}[z;\alpha+1,\beta+1] \qquad (n=1,2,\ldots), \tag{7.4}$$

$$E_n[z;\alpha,\beta] = P_n[z;\alpha,\beta] + \frac{n}{n+\alpha+\beta+1}\,(z-z^{-1})\,P_{n-1}[z;\alpha+1,\beta+1] \qquad (n=0,1,2,\ldots), \tag{7.5}$$

where $nP_{n-1} := 0$ for $n = 0$.

Starting with the operator $Y = Y_{a,b,c,d;q}$ given by (4.4) we can also obtain a differential reflection operator

$$Y = Y_{\alpha,\beta} := \lim_{q\uparrow 1} (1-q)^{-1}\Big(Y_{q^{\alpha+\frac{1}{2}},-q^{\beta+\frac{1}{2}},q^{\frac{1}{2}},-q^{\frac{1}{2}};q} - 1\Big)$$

as a limit case. Then

$$(Yf)[z] = -zf'[z] + \frac{\alpha+\beta+1+(\alpha-\beta)z}{1-z^2}\,(f[z]-f[z^{-1}]) - (\alpha+\beta+1)f[z]. \tag{7.6}$$

Similarly to (4.5) and (4.6) and as a limit case of them, the polynomials (7.4), (7.5) are eigenfunctions of the operator (7.6):

$$YE_{-n} = n\,E_{-n} \qquad (n=1,2,\ldots), \tag{7.7}$$

$$YE_n = -(n+\alpha+\beta+1)\,E_n \qquad (n=0,1,2,\ldots). \tag{7.8}$$

The method of (4.7), (4.9) to rewrite eigenvalue equations for nonsymmetric polynomials as similar equations for vector-valued symmetric polynomials with a matrix-valued operator, also has a limit case here. Now the identifications in (4.9) are based on the rules

$$f[z] = f_1[z] - (z-z^{-1})f_2[z] \quad (f_1, f_2 \text{ symmetric Laurent polynomials}), \tag{7.9}$$

$$(Af)[z] = (A_{11}f_1 + A_{12}f_2)[z] - (z-z^{-1})(A_{21}f_1 + A_{22}f_2)[z]. \tag{7.10}$$

Accordingly, (7.7), (7.8) hold with

$$E_{-n}[z] = (P_n[z;\alpha,\beta), P_{n-1}[z;\alpha+1,\beta+1]) \qquad (n=1,2,\ldots), \tag{7.11}$$

$$E_n[z] = \Big(P_n[z;\alpha,\beta), -\frac{n}{n+\alpha+\beta+1}\,P_{n-1}[z;\alpha+1,\beta+1]\Big) \qquad (n=0,1,2,\ldots) \tag{7.12}$$

and

$$Y = \begin{pmatrix} -(\alpha+\beta+1) & (z^2-1)\frac{d}{dz} + (\alpha+\beta+2)(z+z^{-1}) + 2(\alpha-\beta) \\ (1-z^{-2})^{-1}\frac{d}{dz} & 0 \end{pmatrix}. \tag{7.13}$$

Note that, in view of (7.11), (7.12) and (7.1) the eigenvalue equations (7.7), (7.8) are equivalent to the well-know pair of shift operator relations [10, (1.8.6), (1.8.7)] for Jacobi polynomials.

Remark 7.1. The operator Y in (7.6) coincides, up to a constant term, with Cherednik's [2, line after (3.25)] trigonometric Dunkl operator in the case of root system BC_1. In the notation of [9, (1.2)] this is the operator

$$\frac{d}{dt} - \tfrac{1}{2}(k_1+2k_2) + \left(\frac{k_1}{1-e^{-t}} + \frac{2k_2}{1-e^{-2t}}\right)(1-s),$$

with s the reflection operator. After substitution of $z = e^{-t}$, $k_1 = \alpha - \beta$, $k_2 = \beta + \frac{1}{2}$ this gives $Y + \frac{1}{2}(\alpha+\beta+1)$ with Y as in (7.6). Earlier than [2], Heckman [8] (see also [9, (1.12)]) proposed a trigonometric generalization of the Dunkl operator Dunkl operator which is different from Cherednik's version. For BC_1 this operator becomes

$$\frac{d}{dt} + \tfrac{1}{2}\left(k_1\frac{1+e^{-t}}{1-e^{-t}} + 2k_2\frac{1+e^{-2t}}{1-e^{-2t}}\right)(1-s).$$

After substitution of $t = 2i\theta$ and division by $2i$, and with k_1 and k_2 as before, this gives the operator considered by Chouchene [4, p.1].

In a similar way as before we can prove that the vector-valued polynomials E_n given by (7.4), (7.5) are orthogonal with respect to the symmetric bilinear form

$$\langle g,h\rangle = \langle (g_1,g_2),(h_1,h_2)\rangle = \langle g_1,h_1\rangle_{\alpha,\beta} + \frac{16(\alpha+1)(\beta+1)}{(\alpha+\beta+2)(\alpha+\beta+3)}\,\langle g_2,h_2\rangle_{\alpha+1,\beta+1}\,. \tag{7.14}$$

Here $\langle\,.\,,\,.\,\rangle_{\alpha,\beta}$ is as in (7.2). This form is positive definite if $\alpha,\beta > -1$. The inner product (7.14) can also be written in terms of an integral with positive weight function. First observe that (7.9) implies that

$$f\left[-\left(x \pm i\sqrt{1-x^2}\right)^2\right] = f_1(1-2x^2) \pm 4ix\sqrt{1-x^2}\,f_2(1-2x^2) \qquad (x\in[-1,1]). \tag{7.15}$$

Hence, for Laurent polynomials g,h we obtain from (7.14) and (7.15) that

$$\begin{aligned}\langle g,h\rangle &= \frac{\Gamma(\alpha+\beta+2)}{\Gamma(\alpha+1)\Gamma(\beta+1)} \\ &\quad\times \int_{-1}^{1} g\Big[-\big(x\pm i\sqrt{1-x^2}\big)^2\Big]\,\overline{h\Big[-\big(x\pm i\sqrt{1-x^2}\big)^2\Big]}\,|x|^{2\alpha+1}(1-x^2)^{\beta}\,dx \qquad (7.16)\\ &= \frac{2^{-(2\alpha+2\beta+4)}\Gamma(\alpha+\beta+2)}{\Gamma(\alpha+1)\Gamma(\beta+1)}\int_{-\pi}^{\pi} g(e^{i\theta})\,\overline{h(e^{i\theta})}\,\left|(1-e^{i\theta})^{\alpha+\frac{1}{2}}\,(1+e^{i\theta})^{\beta+\frac{1}{2}}\right|^2 d\theta. \qquad (7.17)\end{aligned}$$

So the orthogonality of the vector-valued polynomials E_n with respect to the inner product (7.14) can be rewritten as the following orthogonality for the Laurent polynomials E_n given by (7.4), (7.5):

$$\int_{-1}^{1} E_m\Big[-\big(x+i\sqrt{1-x^2}\,\big)^2\Big]\,\overline{E_n\Big[-\big(x+i\sqrt{1-x^2}\,\big)^2\Big]}\,|x|^{2\alpha+1}(1-x^2)^{\beta}\,dx=0 \qquad (m\neq n), \tag{7.18}$$

or equivalently,

$$\int_{-\pi}^{\pi} E_m\big[e^{i\theta}\,\big]\,\overline{E_n\big[e^{i\theta}\,\big]}\;\Big|(1-e^{i\theta})^{\alpha+\frac{1}{2}}\,(1+e^{i\theta})^{\beta+\frac{1}{2}}\Big|^2\,d\theta=0 \qquad (m\neq n). \tag{7.19}$$

In the form (7.19) the orthogonality relations for nonsymmetric Jacobi polynomials are the specialization to root system BC_1 of Opdam's orthogonality relations [17, Definition 2.1] for nonsymmetric Jacobi polynomials associated with general root systems. The nonsymmetric Jacobi polynomials which occur as eigenfunctions of Heckman's trigonometric Dunkl operator in case BC_1 are also orthogonal with respect to the inner product (7.17), see Chouchene [4, (2.41)].

Jacobi polynomials can also be obtained as limits of little q-Jacobi polynomials (6.1) (see [10, (5.12.1)]). Define

$$\widetilde{P}_n(x;\alpha,\beta):=\lim_{q\uparrow 1} P_n(x;q^{\alpha},q^{\beta};q). \tag{7.20}$$

These are monic Jacobi polynomials with orthogonality interval rescaled to $[0,1]$:

$$\widetilde{P}_n(x;\alpha,\beta)=\frac{(-1)^n(\alpha+1)_n}{(n+\alpha+\beta+1)_n}\;{}_2F_1\left(\begin{matrix}-n,n+\alpha+\beta+1\\ \alpha+1\end{matrix};x\right). \tag{7.21}$$

By comparing (7.21) with (7.1) we see that

$$P_n[z;\alpha,\beta]=(-1)^n\,2^{2n}\,\widetilde{P}_n\Big(\frac{2-z-z^{-1}}{4};\alpha,\beta\Big). \tag{7.22}$$

Corresponding to (7.20) we can take limits of formulas (6.4)–(6.9) involving nonsymmetric little q-Jacobi polynomials in vector-valued form. Thus we will arrive at formulas which are essentially the same as (7.11)–(7.14) and which can be fully identified with each other by using (7.22). However the expressions (7.4), (7.5) and (7.6) involving Laurent polynomials cannot be obtained as limits of formulas for little q-Jacobi polynomials, since such formulas are missing there.

8 From nonsymmetric Jacobi to nonsymmetric Bessel

In this final section we come back to the nonsymmetric Bessel functions, discussed in section 2, and we show that they are limits of nonsymmetric Jacobi polynomials. There is a well-known

limit from Jacobi polynomials to Bessel functions, see [15, (18.11.5)], which we can rewrite in terms of the notations (7.1) and (2.1) as

$$\lim_{n\to\infty} \frac{2^{\alpha+\beta}\Gamma(\alpha+1)}{\pi^{\frac{1}{2}} n^{\alpha+\frac{1}{2}}} P_n\left(1-\frac{\lambda^2 x^2}{2n^2};\alpha,\beta\right) = \mathcal{J}_\alpha(\lambda x). \tag{8.1}$$

Now from (7.4), (7.5), rewritten in the form of (7.15), we see that (8.1) implies:

$$\lim_{n\to\infty} \frac{2^{\alpha+\beta}\Gamma(\alpha+1)}{\pi^{\frac{1}{2}} n^{\alpha+\frac{1}{2}}} E_{\pm n}\Big[-\Big(\tfrac{\lambda x}{2n} \pm i\sqrt{1-\tfrac{\lambda^2 x^2}{4n^2}}\Big)^2\Big] = \mathcal{J}_\alpha(\lambda x) \pm \frac{i\lambda x}{2(\alpha+1)}\mathcal{J}_{\alpha+1}(\lambda x) = \mathcal{E}_\alpha(\pm\lambda x). \tag{8.2}$$

Corresponding with (8.2) there are limits from (7.6), (7.7), (7.8) to (2.5) and (2.6).

Finally, the limit case of (7.11), (7.12), (7.13) turns down to (2.6) rewritten in vector-valued form as

$$\begin{pmatrix} 0 & x\frac{d}{dx}+2(\alpha+1) \\ \frac{1}{x}\frac{d}{dx} & 0 \end{pmatrix} \begin{pmatrix} \mathcal{J}_\alpha(\lambda x) \\ \frac{i\lambda}{2(\alpha+1)}\mathcal{J}_{\alpha+1}(\lambda x) \end{pmatrix} = i\lambda \begin{pmatrix} \mathcal{J}_\alpha(\lambda x) \\ \frac{i\lambda}{2(\alpha+1)}\mathcal{J}_{\alpha+1}(\lambda x) \end{pmatrix}. \tag{8.3}$$

References

[1] R. Askey and J. Wilson, Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials, *Mem. Amer. Math. Soc.* (1985), no. 319.

[2] I. Cherednik, Double affine Hecke algebras, Knizhnik-Zamolodchikov equations, and Macdonald's operators, *Int. Math. Res. Not.* (1992), no. 9, 171–180.

[3] T. S. Chihara, *An introduction to orthogonal polynomials*, Gordon and Breach, 1978.

[4] F. Chouchene, Harmonic analysis associated with the Jacobi-Dunkl operator on $]-\pi/2,\pi/2[$, *J. Comput. Appl. Math.* 178 (2005), 75–89.

[5] C. F. Dunkl, Hankel transforms associated to finite reflection groups, in *Hypergeometric functions on domains of positivity, Jack polynomials, and applications*, Contemp. Math. 138, Amer. Math. Soc., 1992, pp. 123–138.

[6] C. F. Dunkl and Y. Xu, *Orthogonal polynomials of several variables*, Cambridge University Press, 2001.

[7] G. Gasper and M. Rahman, *Basic hypergeometric series*, 2nd edn., Cambridge University Press, 2004.

[8] G. J. Heckman, An elementary approach to the hypergeometric shift operators of Opdam, *Invent. Math.* 103 (1991), 341–350.

[9] G. J. Heckman, Dunkl operators, in *Séminaire Bourbaki, Vol. 1996/97*, Astérisque 245 (1997), Exp. No. 828, 4, pp. 223–246; `http://www.math.ru.nl/~heckman/Heck_11.ps`.

[10] R. Koekoek and R. F. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue*, Report 98-17, Faculty of Technical Mathematics and Informatics, Delft University of Technology, 1998; `http://aw.twi.tudelft.nl/~koekoek/askey/`.

[11] R. Koekoek, P. A. Lesky and R. F. Swarttouw, *Hypergeometric orthogonal polynomials and their q-analogues*, Springer-Verlag, 2010.

[12] T. H. Koornwinder, Askey-Wilson polynomials as zonal spherical functions on the $SU(2)$ quantum group, *SIAM J. Math. Anal.* 24 (1993), 795–813.

[13] T. H. Koornwinder, The relationship between Zhedanov's algebra AW(3) and the double affine Hecke algebra in the rank one case, *SIGMA* 3 (2007), 063; `arXiv:math/0612730v4`.

[14] I. G. Macdonald, *Affine Hecke algebras and orthogonal polynomials*, Cambridge University Press, 2003.

[15] *NIST Handbook of Mathematical Functions*, Cambridge University Press, 2010; `http://dlmf.nist.gov`.

[16] M. Noumi and J. V. Stokman, Askey-Wilson polynomials: an affine Hecke algebraic approach, in *Laredo Lectures on Orthogonal Polynomials and Special Functions*, Nova Sci. Publ., Hauppauge, NY, 2004, pp. 111–144; `arXiv:math/0001033v1 [math.QA]`.

[17] E. M. Opdam, Harmonic analysis for certain representations of graded Hecke algebras, Acta Math. 175 (1995), 75–121.

[18] M. Rösler and M. Voit, Dunkl theory, convolution algebras, and related Markov processes, in *Harmonic & stochastic analysis of Dunkl processes*, Hermann, Paris, 2008, pp. 1–112; `http://tinyurl.com/29ljycr`.

[19] S. Sahi, Nonsymmetric Koornwinder polynomials and duality, *Ann. of Math.* (2) 150 (1999), 267–282.

[20] S. Sahi, Some properties of Koornwinder polynomials, *Contemporary Math.* 254, Amer. Math. Soc., 2000, pp. 395–411.